

THE UNITED REPUBLIC OF TANZANIA
NATIONAL EXAMINATIONS COUNCIL OF TANZANIA
ADVANCED CERTIFICATE OF SECONDARY EDUCATION
EXAMINATION 2024

132/3A

CHEMISTRY 3A
ACTUAL PRACTICAL A

3 HOURS PRACTICAL ADVANCE INSTRUCTIONS

1.0 IMPORTANT

- 1.1 GREAT CARE MUST BE TAKEN NOT TO DIVULGE THESE INSTRUCTIONS TO BOTH CANDIDATES AND UNAUTHORIZED PERSONS EITHER DIRECTLY OR INDIRECTLY.
- 1.2 MAKE SURE THAT THE CANDIDATES ARE PROVIDED WITH CHEMICALS AND APPARATUSES AS INDICATED IN THESE 3 HOURS PRACTICAL ADVANCE INSTRUCTIONS ONLY AND NOT OTHERWISE.

2.0 PREPARATION AND LABELLING OF CHEMICALS AND APPARATUSES

2.1 Question 1

- Prepare 0.2 M HCl, label it **M2** and allow 100 cm³ per candidate. Follow the following procedure to prepare the solution:

Choose from the following table the appropriate volume of stock (concentrated) solution to be diluted to make 1 litre solution basing on the specification indicated on the bottle of the stock solution.

S/N	Percentage Purity/Assay or its Average when Given in a Range	Density of a Stock Solution (g/cc)	Volume of Stock Solution to be Diluted to make 1 L (cm ³)
1	31 - 32	1.16	19.96
2	≈ 34	1.18	18.18
3	≈ 35	1.18	17.66
4	≈ 36	1.18	17.17
5	≈ 37	1.18	16.70
6	≈ 38	1.18	16.26

- Dissolve 4.24 g anhydrous sodium carbonate and 0.84 g sodium hydrogen carbonate in 1 dm³ of solution and label the resulting mixture as **M1**. Provide each candidate with 100 cm³ of the solution.
- Provide each candidate with a 10 cm³ measuring cylinder.
- Provide the candidates with a 100 cm³ measuring cylinder for sharing in the ratio of 1:4.
- Provide phenolphthalein indicator to each candidate, label it **POP**.

CONFIDENTIAL

- Provide methyl orange indicator to each candidate, label it **MO**.
- Provide each candidate with 1 burette, 1 white tile, 1 pipette (20 cm³ or 25 cm³), 2 titration flasks and 1 retort stand with accessories.
- Provide each candidate with pipette filler.

2.2 Question 2

- Prepare 0.02 M by dissolving 3.16 of KMnO₄ in water to make 1 litre solution and label it **TZ**. Allow 50 cm³ per candidate.
- Dissolve 1.575 g of H₂C₂O₄·2H₂O in 0.25 dm³ of 0.5 M H₂SO₄ and label it **TY**. Allow 50 cm³ per candidate.
- Provide each candidate with a stop watch.
- Provide each candidate with a 250 or 300 cm³ beaker.
- Provide each candidate with two measuring cylinder of 10 cm³.
- Provide each candidate with 2 boiling test tubes.
- Provide each candidate with a thermometer (0 °C – 100 °C).
- Prepare a heat source or burner for sharing in the maximum ratio of 1:4.
- Provide each candidate with 2 test tube holders.
- Provide wire gauze and tripod stand.

2.3 Question 3

- Provide each candidate with 4 g of hydrated ammonium iron(II) sulphate (FeSO₄(NH₄)₂SO₄·6H₂O).
- Provide each candidate with 4 strips of both red and blue litmus papers.
- Provide about 300 cm³ distilled water per candidate.
- Provide each candidate with 4 pyrex test tubes.
- Provide dilute hydrochloric acid, sodium hydroxide, barium chloride, lead ethanoate, ethanoic acid, concentrated sulphuric and hydrochloric acid, potassium hexacyanoferrate(III) as bench reagents.
- Prepare heat source or burner for sharing in the maximum ratio of 1:4.