

L

**THE UNITED REPUBLIC OF TANZANIA
NATIONAL EXAMINATIONS COUNCIL
ADVANCED CERTIFICATE OF SECONDARY EDUCATION
EXAMINATION**

131/3A

**PHYSICS 3A
ACTUAL PRACTICAL A
(For Both School and Private Candidates)**

Time: 3:10 Hours

Monday, 18th February 2013 a.m.

Instructions

1. This paper consists of **three (3)** questions.
2. Answer all questions.
3. Question **Number 1** carries 20 marks and the other **two (2)**, 15 marks each.
4. Calculations should be clearly shown.
5. Mathematical tables and non-programmable calculators may be used.
6. Cellular phones are **not** allowed in the examination room.
7. Write your **Examination Number** on every page of your answer booklet (s).
8. Use the following:

$$\pi = 3.14$$



1. The aim of this experiment is to determine the acceleration due to gravity, g , by simple pendulum.

Proceed as follow:




Figure 1

- Using a pointer, place a reference mark at the equilibrium position of the bob. Set the bob to oscillate through a small angle and begin counting the number of oscillations through the equilibrium position of the bob. Find the time for 30 complete oscillations.
- Measure the length, l , of the thread from the point of suspension to the point of attachment of the bob and record the result in metres (m).
- By raising the thread each time, shorten the length of the pendulum by about 10cm, 20cm, 30cm and 40cm.
- On each occasion, find the length of the thread in metres and the time for 30 complete oscillations. Tabulate your results.
- Plot the graph of l against T^2 , where T is the period of the oscillations related by the equation:

$$T = 2\pi \sqrt{\frac{l+c}{g}}$$

and c is the small constant length from the point of attachment of the bob to the centre of gravity of the bob.

- Determine the acceleration due to gravity, g .
 - Mention any two sources of errors.
2. You are provided with a calorimeter and stirrer, thermometer, water bath containing hot water maintained at 90°C , stopwatch and a cardboard.

Proceed as follows:

- Fill the calorimeter with hot water about $\frac{3}{4}$ full. Put the calorimeter containing

hot water on a cardboard placed over the bench. Starting with $\theta = 75^{\circ}\text{C}$ record temperature after 2 minutes intervals. Take your reading for 20 minutes.

Note: Stir water thoroughly until the end of your experiment.

- (b) Record the room temperature at the beginning and at the end of the experiment. Find the average temperature $\theta_s(^{\circ}\text{C})$.
- (c) Tabulate your results.
- (d) Plot a graph of $\log_{10}(\theta - \theta_s)$ against time, t .
- (e) Given that $A = \log_{10}(\theta - \theta_s) + Kt$ where A is a constant.

Determine the value of K and constant A .

- (f)
 - (i) From the graph, state the relation between the rate of loss of heat from a calorimeter and the excess temperature over its surrounding.
 - (ii) Which law does this relation obey?

3. The aim of this experiment is to determine the resistivity ' ρ ' of the wire labeled X and the internal resistance of the battery provided.

Proceed as follows:




Figure 2

- (a) Connect the circuit as shown in Figure 2. With the plug key open adjust the length of wire X to a value of $L_x = 20\text{cm}$. Note the ammeter reading.

Note: The plug key should remain open throughout the experiment.

- (b) Repeat the procedures in part (a) above for $L_x = 40\text{cm}$, 60cm , 80cm and 100cm each time recording the ammeter reading.
- (c) Tabulate your results as shown in Table 1.

Table 1

Length L_x of wire X (cm)	Current I (A)	$\frac{1}{I}$ (A^{-1})

- (d) Plot a graph of $\frac{1}{I}$ against L_x .
- (e) Determine the slope and the $\frac{1}{I}$ intercept of the graph.
- (f) Measure and record the diameter of a wire.
- (g) Using your graph, determine the resistivity ρ of the wire and internal resistance r of the battery.