## THE UNITED REPUBLIC OF TANZANIA NATIONAL EXAMINATIONS COUNCIL CERTIFICATE OF SECONDARY EDUCATION EXAMINATION

032/2C

## CHEMISTRY 2C ACTUAL PRACTICAL C

(For Both School and Private Candidates)

Time: 2:30 Hours

Wednesday, 19th November 2014 a.m.

## Instructions

- This paper consists of three (3) questions. Answer all the questions.
- Question 1 carries twenty (20) marks and the rest carry fifteen (15) marks each.
- Qualitative Analysis Guidance Pamphlets may be used after a thorough check by the supervisor.
- 4. Cellular phones and calculators are not allowed in the examination room.
- 5. Write your **Examination Number** on every page of your answer booklet(s).
- 6. You may use the following constants:

Atomic masses:

H = 1, 
$$C = 12$$
,  $O = 16$ ,  $Na = 23$ ,  $Cl = 35.5$ ,  $S = 32$ .  
1 litre = 1 dm<sup>3</sup> = 1000 cm<sup>3</sup>.

- You are provided with the following solutions: 1.
  - D: Containing 1.46 g of hydrochloric acid in 0.40 dm<sup>3</sup> of solution; D: Containing 1.46 g of hydrochloric acid in 0.40 day.

    F: Containing 2.0 g of impure sodium hydroxide contaminated with sodium chloride.
  - 0.5 dm<sup>3</sup>:

Phenolphthalein and methyl orange indicators.

**Ouestions** 

- Which is the suitable indicator for the titration of the given solutions? Give (a) reason for your answer.
- Titrate the acid (in a burette) against the base (in a conical flask) using drops of your indicator and obtain three titre values.
- cm<sup>3</sup> of F required \_\_\_\_ cm<sup>3</sup> of D for complete reaction. Write a balanced chemical equation for the reaction between D and F. (c) (ii)
- Showing your procedures clearly, calculate the percentage by mass of sodius (d) chloride in impure base.
- You are provided with the following: 2.

Solution C containing 0.5 M sodium thiosulphate (Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>);

Solution E containing 0.1 M nitric acid (HNO<sub>3</sub>);

Distilled water;

Plain paper marked X;

Stopwatch.

## **Procedure**

Measure 10 cm<sup>3</sup> of solution C and put into 100 cm<sup>3</sup> beaker provided.

(ii) Measure 10 cm<sup>3</sup> of solution E and put into 100 cm<sup>3</sup> beaker containing solution C, and immediately start the stopwatch.

- (iii) Swirl the contents and place the beaker on top of letter X on the plain paper provided Watch from above and observe the changes.
- (iv) Switch off the stopwatch when the mark X disappears.
- (v) Record the time taken for the letter X to disappear.
- (vi) Repeat the experiment using different data as shown in Table 1.

Table 1: Table of Results

Experiment	Vol. of E (cm <sup>3</sup> )	Vol. of C (cm <sup>3</sup> )	Vol. of Distilled water (cm <sup>3</sup> )	Time (s)
1	10	10	0	
2	10	8	2	
3	10	6	4	
4	10	4	6	

Oue	estions
(a)	What is the aim of the whole experiment?
(b)	Complete Table 1.
(c)	Giving reason(s), identify the experiment in

- n which the reaction was:
  - (1) fast
  - (ii) slow.
- With state symbols, write the ionic equation for the reaction between C and E. (d)
- List any three factors affecting the reaction in 2(d). (e)
- Plot a graph of volume of C against time. (i) (f)
  - What can you conclude from the graph? (ii)

Sample Y contains one cation and one anion. Using systematic qualitative analysis procedures record carefully your experiments, observations, inferences and finally identify the anion and cation in sample Y.

Table 2: Table of results

S/n	Experiment	Observation	Inference

-		فسنتانا	•	
ľ	nc	110	ion	
$\mathbf{v}$	110	us.	IUII	

(i)	The estion	:1	- 37		
(1)	The cation	ın sampı	e r	18	

(ii)	The anion	in	sample	Y	is	 •
			1			

<sup>(</sup>iii) The chemical formula of sample Y is \_\_\_\_\_.