DIPLOMA IN SECONDARY EDUCATION EXAMINATION

732/2A

(ALTERNATIVE A PRACTICAL)

Time: 3 Hours

Tuesday, 18th May 2010 a.m.

Instructions

- 1/22-2/This paper consists of three questions.
- 2//cc Answer all the questions
- 3//65-3/Question humber one carries 40/marks and question two and three carries 30 marks and question two and three carries 30 marks and question two and three carries 30/0/22-30/0/2
- 4. You are allowed to use qualitative analysis guide pamphlets for answering question number 3.
- 5. Cellular phones are **not** allowed in the examination room.
- 6. Write your Examination Number on every page of your answer booklet(s).
- 7. The following constants might be useful in your calculations:

Atomic mass:

Na = 23, O = 16, H = 1, C = 12, K = 39, Mn = 55, S = 32

1. You are provided with the following:

T1: A solution containing 40 g of Na₂S₂O_{3.5}H₂O dm⁻³.

A solution of dilute HCl

T3: Distilled water A stop watch.

Procedure:

- Using blue or black pen write a clear letter x on a piece of white paper and place a small beaker on top of the letter x such that the letter x is visible (a) through the solution.
- Put 10 cm³ of T1 into the beaker followed by 5 cm³ of T2 and start the stop watch. Stir gently and record the time taken for the letter x to disappear in (b) the solution.
- Repeat the above procedures by using various amounts of T1 and distilled (c) water T3 as indicated in the table below.

Table of Results:

	Table of Re					7-7	$\frac{1}{-(\sec^-)}$
	Volume of T1 (cm ³)	Volume of H ₂ O (cm ³)	Volume of T2 (cm ³)	$\frac{[S_2O_3]}{M}$	t(sec)	$\left[S_2O_3^{2-}\right]^{x t} (se$	c) -(sec)
ŀ	10	0	5				
1	8	2	5	2 3 3 3 5 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1			
r	6	4	5				4
\mid	4	6	5				

- Complete the table above. (d)
- Plot a graph of $\left[S_2O_3^{2-}\right]$ against t. (e)
- Plot a graph of $\frac{1}{t}$ against $\left[S_2O_3^{2-1}\right]$. (f)
- From your graphs what is the effect of $\left[Na_2S_2O_3\right]$ on the rate of reaction? (g)
- What is the order of reaction with respect to $[Na_2S_2O_3]$? (h)

- 2. You are provided with the following solutions:
 - A solution of 5.032 g of dibasic organic acid of anhydrous salt with molecular mass 90 made to one litre of aqueous solution.
 - B: A solution of 0.1 M NaOH
 - Phenolphthalein indicator. **C**:

Procedure:

- Pipette 20 cm³ or 25 cm³ of A into a conical flask. (a)
- Add to it few drops of C. (b)
- Put B into the burette. (c)
- Titrate B against A until the colour change is observed. (d)

Table of results:

Tit					
Titration Number		Pilot	1	2	3
Final volume (cm³)					1
Initial volume (cm ³)					
Volume used (cm ³)		3	6	1) G**

- (a) The volume of pipette used was cm³
- (b) The volume of burette was cm³ For complete neutralization _____cm³ of A required ____cm³ of solution B
- Write a balanced equation for the reaction taking place between A and B. (c)
- Calculate the (d)
 - molarity of A. (i)
 - concentration of A in gdm⁻³. (ii)
 - (iii) number of moles of water of crystallization per mole of hydrated dibasic organic acid.
- Sample W is a compound containing one anion and one cation. Using qualitative analysis techniques, identify the two ions and hence deduce the name of the 3. compound.