# THE UNITED REPUBLIC OF TANZANIA NATIONAL EXAMINATIONS COUNCIL DIPLOMA IN SECONDARY EDUCATION EXAMINATION

731/2A

## PHYSICS 2A (ACTUAL PRACTICAL A)

Time: 3 Hours

Thursday, 10<sup>th</sup> May 2012 a.m.


## Instructions

- 1. This paper consists of three (3) questions.
- 2. Answer all questions.
- 3. Question number one carries 40 marks, question number two and three carries 30 marks each.
- 4. Mathematical tables and non-programmable calculators may be used.
- 5. Cellular phones are **not** allowed in the examination room.
- 6. Write your **Examination Number** on every page of your answer booklet(s).

The aim of this experiment is to determine the force constant of the spring k.

#### Apparatus:

You are provided with a weight pan, meter rule, pointer, spiral spring, slotted weights, stop watch and two retort stands with clamps.



#### Figure 1

#### **Procedures:**

- (a) Set up the apparatus provided for this experiment as shown in Figure 1 above.
- (b) Record the scale reading  $s_0$ .
- (c) Add 50 gm on the weight pan and record the new scale reading s.
- (d) Calculate the extension  $(e = s s_0)$  caused by the weight.
- (e) Repeat with different load of 100 gm, 150 gm, 200 gm, 250 gm, until 600 gm.

# **Questions:**

- (i) Tabulate your results.
- (ii) Sketch a well labeled diagram of the experiment.
- (iii) Plot a graph of load against extension.
- (iv) Find the gradient G of your graph.
- (v) What is the physical meaning of the gradient?
- (vi) State four sources of errors and suggest the ways to minimize.

Page 2 of 4

You are provided with a thermometer, a calorimeter, a stirrer, a stopwatch/stop clock, beaker and a liquid labeled B.

#### **Procedures:**


- (a) Using a beaker, take about 200cm<sup>3</sup> of the liquid B and heat until it boils.
- (b) Quickly transfer the boiling liquid B from a beaker to the calorimeter provided. Record the temperature and immediately start the watch.
- (c) While stirring the liquid and constantly fanning with the piece of paper provided, note and record the temperature of the liquid as it cools at intervals of 2 minutes. Continue with reading the temperature for 20 minutes.

### **Questions:**

(i) Tabulate your results for  $\theta$  and t, where  $\theta$  is the temperature in °C and t is the time in minutes.

_									.1195	Evaluate r for one	(IV)
Ľ	Time t (Minute)						1.			State the source of	daily i
Ľ	Femperature $\theta$ °C	٠.	ì	•	•	1. 1	•	: '			

- (ii) Draw a well labeled diagram of the experiment.
- (iii) Plot a graph of  $\theta$  against t.
- (iv) Determine the rate of cooling of liquid B at  $70^{\circ}C_{1}$   $60^{\circ}C_{1}$   $50^{\circ}C_{2}$
- (v) Mention four sources of error in this experiment.
- 3. The aim of this experiment is to determine the electromotive force (E) and internal resistance (r) of a cell.



## **Procedures:**

- (a) Connect the circuit as shown in Figure 2 above.
- (b) Put  $R=1\Omega$  and quickly read the value i on the ammeter.
- (c) Repeat the procedure in 3(a) above for values of  $R=2\Omega$ ,  $3\Omega$ ,  $4\Omega$ ,  $5\Omega$  and  $6\Omega$ .

Page 3 of 4

## **Questions:**

will invested

(i) Tabulate your results and complete the following table

Resistance $R(\Omega)$	Current i(A)	$\frac{1}{i}(\mathbf{A}^{-1})$
1		
2		
3	ert di liberio de 1808 il	g II to appet the
4		
5		
6	of the late of the second of the second	charles !

- (ii) Plot the graph R against  $\frac{1}{i}$ .
  - (iii) Using the graph and the equation  $R = \frac{E}{i} r$ . Find the value of E.
  - (iv) Suggest how E and r may be evaluated from your graph.
  - (v) Evaluate E for one cell.
  - (vi) Evaluate r for one cell.

e (E) and internal toxidanc

(vii) State the source of error and suggest one way of minimizing it.